


## LARINGE

Anni 2005-2017. Sintesi dei dati di incidenza. LARINGE

	MASCHI	FEMMINE
<b>Numero casi</b>	386	28
<b>% sul totale</b>	1,4	0,1
<b>Tasso grezzo</b>	10,4	0,7
<b>Tasso standardizzato</b>	10,8	0,7
<b>Rischio cumulativo</b>	0,7	0
<b>Età media</b>	66,9	65,3
<b>Età mediana</b>	66	65


Anni 2005-2017. Trend dei dati di incidenza. LARINGE

	MASCHI			FEMMINE		
	Num_M	TSD_M	ES_TSD_M	Num_F	TSD_F	ES_TSD_F
<b>2005</b>	28	11,37	2,19	1	0,43	0,43
<b>2006</b>	25	9,69	1,96	1	0,23	0,23
<b>2007</b>	34	12,69	2,19	1	0,30	0,30
<b>2008</b>	22	8,87	1,92	2	0,60	0,43
<b>2009</b>	26	9,89	1,95	2	0,59	0,42
<b>2010</b>	29	10,61	1,98	1	0,35	0,35
<b>2011</b>	39	13,56	2,18	3	1,18	0,69
<b>2012</b>	31	10,83	1,95	2	0,53	0,39
<b>2013</b>	38	13,54	2,21	5	1,59	0,72
<b>2014</b>	31	11,24	2,03	3	0,98	0,57
<b>2015</b>	28	9,69	1,84	3	1,01	0,59
<b>2016</b>	31	10,88	1,98	0	0,00	0,00
<b>2017</b>	24	8,56	1,76	4	1,27	0,64




Anni 2005-2017. Tassi età specifici di incidenza LARINGE

ETA'	MASCHI	FEMMINE
0-04	0,00	0,00
05-09	0,00	0,00
10-14	0,00	0,00
15-19	0,00	0,00
20-24	0,00	0,00
25-29	0,00	0,00
30-34	0,39	0,00
35-39	0,75	0,00
40-44	1,42	0,35
45-49	6,42	0,70
50-54	9,88	0,74
55-59	24,00	1,21
60-64	31,31	1,84
65-69	22,19	2,46
70-74	36,83	2,53
75-79	37,25	1,56
80-84	29,40	1,96
85+	30,20	0,00




## Anni 2005-2017. Sopravvivenza. LARINGE


	2005-09	2010-14	stima 2015-17
1a_M	91,78	84,70	90,00
2a_M	82,62	72,54	87,60
3a_M	70,61	65,52	80,23
4a_M	65,35	64,05	77,17
5a_M	65,84	62,75	73,41
1a_F	43,68	101,47	86,06
2a_F	29,95	81,00	86,37
3a_F	30,32	82,17	86,55
4a_F	30,77	68,26	55,08
5a_F	31,27	69,14	55,08




Anno 2017. Prevalenza osservata. LARINGE

Anni	MASCHI		FEMMINE	
	Casi	Tassi	Casi	Tassi
2009	47	16,78	4	1,38
2010	71	25,34	6	2,07
2011	93	33,19	8	2,76
2012	110	39,26	11	3,79
2013	126	44,97	12	4,14
2014	142	50,83	15	5,17
2015	157	56,19	16	5,51
2016	163	58,33	16	5,51
2017	169	60,47	17	5,86





MASCHI



FEMMINE

